


LIMITS INVOLVING INFINITY

We have already seen vertical asymptotes which occur when

$$\lim_{x \rightarrow a^+} f(x) = +\infty, -\infty$$

$$\lim_{x \rightarrow a^-} f(x) = +\infty, -\infty$$

as in $f(x) = \frac{1}{x-5} + 2$ which has an asymptote at $x=5$:



Notice there is another asymptote at $y=2$ — a horizontal asymptote.

* We use a strategy: ~~divide~~ isolate dominant term.

(2.)

Defⁿ Horizontal Asymptote (HA)


$f(x)$ has the horizontal asymptote $y=b$ when

$$\lim_{x \rightarrow \infty} f(x) = b \quad \text{or} \quad \lim_{x \rightarrow -\infty} f(x) = b.$$

Note a function ~~have~~ can have up to two HAs — one for each infinity: $+\infty, -\infty$.

EXAMPLE

$$y = \frac{x^2 - 1}{x^2 + 1} = f(x)$$



Geometry suggests $\lim_{x \rightarrow \infty} f(x) = 1$.

Algebraically: (using limit laws) *

$$\lim_{x \rightarrow \infty} \frac{x^2 - 1}{x^2 + 1} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x^2}(x^2 - 1)}{\frac{1}{x^2}(x^2 + 1)} = \lim_{x \rightarrow \infty} \frac{\left(\frac{x^2}{x^2} - \frac{1}{x^2}\right)}{\left(\frac{x^2}{x^2} + \frac{1}{x^2}\right)}$$

$$= \lim_{x \rightarrow \infty} \frac{1 - \frac{1}{x^2}}{1 + \frac{1}{x^2}} = \frac{\lim_{x \rightarrow \infty} 1 - \lim_{x \rightarrow \infty} \frac{1}{x^2}}{\lim_{x \rightarrow \infty} 1 + \lim_{x \rightarrow \infty} \frac{1}{x^2}} = \frac{1 - 0}{1 + 0} = 1$$

$$\lim_{x \rightarrow \infty} 1 + \lim_{x \rightarrow \infty} \frac{1}{x^2} \quad \text{STUCK}$$

we have no rule for this

Propⁿ $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, $\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$

Proof: Requires strict defn - omitted for MAT134.

Thus $\lim_{x \rightarrow \infty} \frac{1}{x^2} = \lim_{x \rightarrow \infty} \left(\frac{1}{x}\right) \left(\frac{1}{x}\right) = \lim_{x \rightarrow \infty} \frac{1}{x} \lim_{x \rightarrow \infty} \frac{1}{x} \neq 0$
 $= 0 \cdot 0 = 0.$

and so AHA not stuck anymore.

$\otimes = \frac{1-0}{1+0} = 1 = \lim_{x \rightarrow \infty} \frac{x^2-1}{x^2+1}$

Confirming / Demonstrating / Proving what the geometric intuition suggested.

Defⁿ $\lim_{x \rightarrow \infty} f(x) = L$ (L finite or $L < \infty$) when

there is some point on the x -axis after which $f(x)$ is arbitrarily close to L .



STRICTLY: $\forall \epsilon > 0 \exists a \in \mathbb{R} : x > a \Rightarrow f(x) \in (L - \epsilon, L + \epsilon)$

Defⁿ $\lim_{x \rightarrow -\infty} f(x) = L$ "... before which ..."

$$\forall \epsilon > 0 \exists a \in \mathbb{R}: x < a \Rightarrow |f(x) - L| < \epsilon$$

equivalent to $f(x) \in (L - \epsilon, L + \epsilon)$
(just for the sake of it)

EXAMPLE Functions can cross the HA: ~~an~~ For instance the following functions have HA, despite crossing them.



(ISOLATING DOMINANT TERM)

EXAMPLE $\lim_{x \rightarrow \infty} \frac{3x^2 - x - 2}{5x^2 + 4x + 1} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x^2}(3x^2 - x - 2)}{\frac{1}{x^2}(5x^2 + 4x + 1)}$

$$= \lim_{x \rightarrow \infty} \frac{\left(\frac{3x^2}{x^2} - \frac{x}{x^2} - \frac{2}{x^2} \right)}{\left(\frac{5x^2}{x^2} + \frac{4x}{x^2} + \frac{1}{x^2} \right)} = \lim_{x \rightarrow \infty} \frac{\left(3 - \frac{1}{x} - \frac{2}{x^2} \right)}{\left(5 + \frac{4}{x} + \frac{1}{x^2} \right)}$$

$$= \frac{3 - 0 - 0}{5 + 0 + 0} = 3/5. \quad \text{Check w/ DESMOS.}$$

EXAMPLE : by conjugation and isolating dom term.

$$\lim_{x \rightarrow \infty} \sqrt{x^2+1} - x \quad " = \infty - \infty "$$

$$= \lim_{x \rightarrow \infty} \left[(x^2+1)^{\frac{1}{2}} - x \right] \left[\frac{(x^2+1)^{\frac{1}{2}} + x}{(x^2+1)^{\frac{1}{2}} + x} \right]$$

$$= \lim_{x \rightarrow \infty} \frac{(x^2+1) - x^2}{(x^2+1)^{\frac{1}{2}} + x}$$

$$= \lim_{x \rightarrow \infty} \frac{1}{(x^2+1)^{\frac{1}{2}} + x}$$

Dominant term is $\sqrt{x^2} = x$. so...

$$= \lim_{x \rightarrow \infty} \frac{\frac{1}{x} \cdot 1}{\frac{1}{x} \left[(x^2+1)^{\frac{1}{2}} + x \right]}$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{\left(\frac{x^2}{x^2} + \frac{1}{x^2} \right)^{\frac{1}{2}} + \frac{x}{x}}$$

$$= \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{\left(1 + \frac{1}{x^2} \right)^{\frac{1}{2}} + 1}$$

$$= \frac{0}{(1+0)^{\frac{1}{2}} + 1} = 0/2 = 0.$$

Confirm w/ DESMOS.

$$\sqrt{x^2} = |x|$$

(6)

EXAMPLE: $f(x) = \frac{(2x^2+1)^{\frac{1}{2}}}{3x-5}$ has three asymptotes.

$$\lim_{x \rightarrow \infty} \frac{(2x^2+1)^{\frac{1}{2}}}{3x-5} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x} (2x^2+1)^{\frac{1}{2}}}{\frac{1}{x} (3x-5)}$$

because $x \rightarrow \infty$, $x > 0$ and $\sqrt{x^2} = |x| = x$

$$= \lim_{x \rightarrow \infty} \frac{\frac{1}{\sqrt{x^2}} \sqrt{2x^2+1}}{\frac{1}{x} (3x-5)} = \lim_{x \rightarrow \infty} \frac{\sqrt{\frac{2x^2}{x^2} + \frac{1}{x^2}}}{\left(\frac{3x}{x} - \frac{5}{x}\right)}$$

$$= \lim_{x \rightarrow \infty} \frac{\sqrt{2 + \frac{1}{x^2}}}{\left(3 - \frac{5}{x}\right)} = \lim_{x \rightarrow \infty} \frac{\sqrt{2+0}}{(3-0)} = \sqrt{2}/3.$$

Thus $x = \sqrt{2}/3$ is a horizontal asymptote.

$$\lim_{x \rightarrow -\infty} \frac{(2x^2+1)^{\frac{1}{2}}}{3x-5} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x} (2x^2+1)^{\frac{1}{2}}}{\frac{1}{x} (3x-5)}$$

because $x \rightarrow -\infty$, $x < 0$ and $\sqrt{x^2} = |x| = -x$

$$\lim_{x \rightarrow -\infty} \frac{(2x^2+1)^{\frac{1}{2}}}{3x-5} = \lim_{x \rightarrow \infty} \frac{\frac{-1}{(x^2)^{\frac{1}{2}}} (2x^2+1)^{\frac{1}{2}}}{\frac{1}{x} (3x-5)} = \dots = -\sqrt{2}/3.$$

So there is a HA at $x = -\sqrt{2}/3$.

The last, vertical asymptote is at $x = 5$ as

$$\lim_{x \rightarrow 5^+} f(x) = +\infty. \quad \lim_{x \rightarrow 5^-} f(x) = -\infty.$$

Limits at Infinity

Note = $\infty + \infty = \infty$ but $\infty - \infty$ undefined.
 $\infty \cdot \infty = \infty$ ∞ / ∞ undefined.

NOTATION $\lim_{x \rightarrow \infty} f(x) = \infty$ means $f(x)$ is "growing unbounded" as $x \rightarrow \infty$.

(The only alternative is having a HA).

STRICT $\lim_{x \rightarrow \infty} f(x) = \infty \Leftrightarrow \forall y \exists a: x > a \Rightarrow f(x) > y$.

(i.e. you cannot pick y large enough to "bound" $f(x)$).

There are similar defs for $\lim_{x \rightarrow \pm\infty} f(x) = \pm\infty$

EXAMPLE $\lim_{x \rightarrow \infty} x^2 - x$

WRONG: $\lim_{x \rightarrow \infty} x^2 - x = \lim_{x \rightarrow \infty} x^2 - \lim_{x \rightarrow \infty} x = \infty - \infty = 0$
not true

CORRECT: $\lim_{x \rightarrow \infty} x^2 - x = \lim_{x \rightarrow \infty} x(x-1) = \lim_{x \rightarrow \infty} x \cdot \lim_{x \rightarrow \infty} (x-1)$
 $= \infty \cdot \infty = \infty.$


EXERCISE $\lim_{x \rightarrow \infty} \frac{3x^2 - x + 4}{2x^2 + 5x - 8}$

EXERCISE


• $\lim_{x \rightarrow a} \left(\frac{x^2}{2} - \frac{1}{x} \right)$ for $a = 0^+, 0^-, 2^{3/2}$, and -1 .

§ CONTINUITY

Basically: a function is continuous over some interval when you can draw it w/out lifting your pen.



"Continuous on (a,b)"



"discontinuous at c"


Defⁿ $f: \mathbb{R} \rightarrow \mathbb{R}$ a function is continuous at a when $\lim_{x \rightarrow a} f(x) = f(a)$.

Consequently, to ~~prove~~ show a function is cont. at a you must show three things:

(1) $a \in \text{dom } f$, (2) $\lim_{x \rightarrow a} f(x)$ exists, and

(3) $\lim_{x \rightarrow a} f(x) = f(a)$.

EXAMPLE




$h(x)$ is discont at $x = a, b, c.$

$x = a$ because $a \notin \text{dom } f \Rightarrow \text{FAILS } (1)$

$x = b$ because $\lim_{x \rightarrow b^-} h = 2 \neq 4 = \lim_{x \rightarrow b^+} \Rightarrow \text{FAILS } (2)$

$x = c$ because $\lim_{x \rightarrow c} h = 2 \neq 4 = h(c) \Rightarrow \text{FAILS } (3)$

EXAMPLE $f(x) = \frac{x^2 - x - 2}{x - 2}$ cannot be cont. at $x = 2$ because $2 \notin \text{dom } f.$



Note, we can put the point back in. Because

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \frac{(x+1)(x-2)}{(x-2)} = \lim_{x \rightarrow 2} x+1 = 3.$$


$$g(x) = \begin{cases} 3 & x = 2 \\ \frac{(x+1)(x-2)}{x-2} & \text{elsewhere} \end{cases}$$

Restores continuity at this point - a ~~jump disc.~~ removable disc.

EXAMPLE $f(x) = \lfloor x \rfloor$ "floor" or "truncation"

e.g. $\lfloor 2.3 \rfloor = 2$ $\lfloor 2 \rfloor = 2$


Let $x \geq 0$ then



notice

and so $f(x)$ is disc at every integer.

EXAMPLE



f . Here we say f is left-continuous at $x=a$, but not right-continuous

Defⁿ Right Continuous / Left Continuous

A function f is continuous from the right at $x=a$

when $\lim_{x \rightarrow a^+} f(x) = f(a)$

and left-continuous when

$\lim_{x \rightarrow a^-} f(x) = f(a)$.

QUESTION f cont at $a \Rightarrow f$ L-cont & R-cont at a ?
YES!

Defⁿ f is cont on an open-interval $(a,b) \subseteq \mathbb{R}$
 when $\forall x \in (a,b); f(x)$ is cont.

Moreover, if $f(x)$ is R-cont at $x=a$ and
 L-cont at $x=b$ then $f(x)$ is continuous on
 $[a,b]$.

THM
~~EXAMPLES~~ When f and g are cont at a , $c \in \mathbb{R}$
 then the following are also cont. at a .

① $f+g$ ③ f/g provided $g(a) \neq 0$

② $f \cdot g$ ④ $c \cdot f$ and $c \cdot g$.

Proof: Omitted.

Recall $\mathbb{R}[x] = \{a_0 + a_1x + a_2x^2 + \dots + a_nx^n : a_k \in \mathbb{R}\}$
 are polynomials and ~~a~~

$\{f/g : f, g \in \mathbb{R}[x] \text{ and } g \neq 0\}$

are rational functions.

Thm Polynomials are everywhere cont.

Rational functions are cont. on their domain.

EXAMPLE $\frac{x^3 + 2x - 1}{5 - 3x}$ is continuous everywhere but $x = \frac{5}{3}$.

Thm g cont at a and f cont at $g(a)$
then $(f \circ g)(x)$ is cont at a .

§ Intermediate Value Thm

Suppose f is cont on (a, b) and $f(a) < 0$
and $f(b) > 0$. Given that we cannot pick
up our pens when drawing, $f(x)$ must
pass through the $y=0$ axis.

In other words $\exists c \in (a, b) : f(c) = 0$.


Thm IVT

Provided:

- f is cont on $[a, b]$
- $N \in [f(a), f(b)]$

Then: $\exists c \in (a, b) : f(c) = N$.

In other words: You must cross $y=N$ to connect p and q below.



Show

EXAMPLE: $f(x) = 4x^3 - 6x^2 + 3x - 2 = 0$ has a root in $(1, 2)$.

Notice • $f(x)$ is a polynomial \Rightarrow everywhere cont

• $f(1) = 4 - 6 + 3 - 2 = -1 < 0$

• $f(2) = 4 \cdot 8 - 6 \cdot 4 + 3 \cdot 2 - 2 = 12 > 0$

⑦

Consequently,

We have $0 \in (f(1), f(2))$ and by NT

$$\exists c \in (1, 2) : f(c) = 0$$

Thus f has a root in $(1, 2)$.